


ORIGINAL ARTICLE

Epidemiology of Allergic Disease

Patterns of allergic sensitization and atopic dermatitis from 1 to 3 years: Effects on allergic diseases

C. Dharma¹ | D. L. Lefebvre¹ | M. M. Tran¹ | W. Y. W. Lou² | P. Subbarao³ |
A. B. Becker⁴ | P. J. Mandhane⁵ | S. E. Turvey⁶ | M. R. Sears¹  | the CHILD Study
investigators^a

¹Department of Medicine, McMaster University, Hamilton, Canada

²Dalla Lana School of Public Health, University of Toronto, Toronto, Canada

³Department of Pediatrics, University of Toronto & Hospital for Sick Children, Toronto, Canada

⁴Department of Pediatrics & Child Health, University of Manitoba, Winnipeg, Canada

⁵Department of Pediatrics, University of Alberta, Edmonton, Canada

⁶Department of Pediatrics, University of British Columbia, Vancouver, Canada

Correspondence

M. R. Sears, Firestone Institute for Respiratory Health, St. Joseph's Healthcare and McMaster University, Hamilton, Ontario, Canada.

Email: searsm@mcmaster.ca

Funding information

Canadian Institutes of Health Research (CIHR), Allergy, Genes and Environment Network of Centres of Excellence (AllerGen NCE, Inc.)

Summary

Background: While allergic sensitization and atopic dermatitis (AD) are known to increase the risk for allergic diseases, the impact of different temporal and clinical patterns of sensitization and AD is less well defined.

Objective: We investigated patterns of sensitization and AD from early infancy to age 3, and the differential risk of developing allergic diseases within each pattern in a general cohort.

Methods: Children (n = 2629) from the Canadian Healthy Infant Longitudinal Development (CHILD) Study underwent skin prick tests and were assessed clinically for AD at ages 1 and 3 years. We applied an unsupervised latent class analysis (LCA) to the following 5 factors at these ages: AD, food sensitization, inhalant sensitization, poly-sensitization to foods and poly-sensitization to inhalants. The risks for developing asthma, allergic rhinitis and food allergy at 3 years were evaluated for each identified group.

Results: Five distinct classes were revealed by LCA: healthy (81.8%), atopic dermatitis (7.6%), inhalant sensitization (3.5%), transient sensitization (4.1%) and persistent sensitization (3.2%). Using healthy children as the baseline, children in the "atopic dermatitis" group had the next lowest risk for all allergic outcomes at 3 years; those in the "inhalant sensitization" group had the highest risk for allergic rhinitis; children in the "transient sensitization" group were at an increased risk for food allergy; while children in the "persistent sensitization" group had the highest risk for all allergic diseases.

Conclusion and Clinical Relevance: There is substantial heterogeneity among allergen-sensitized children. Researchers and clinicians need to be aware of the non-specificity associated with labelling children simply as "atopic" and "non-atopic" without considering the timing of their atopic history, type of sensitization and AD status. Children with AD who were poly-sensitized to foods at an early age appear to be at greatest risk of developing other allergic diseases.

KEYWORDS

allergic rhinitis, asthma, atopic dermatitis, food sensitization, inhalant sensitization

^aCanadian Healthy Infant Longitudinal Development Study: See Appendix 1

1 | INTRODUCTION

Understanding the temporal and clinical patterns of allergic sensitization and atopic dermatitis (AD) from a young age is crucial in identifying children most vulnerable to serious allergic diseases at later ages. Previous prospective cohorts have illustrated that from birth to age one, children are usually only mono-sensitized (sensitized to one allergen), primarily to food. Between ages one to four years, inhalant sensitization becomes more frequent, concurrent with a decline in food sensitization^{1–3}; more children also start to exhibit poly-sensitization (sensitization to multiple allergens).^{4,5}

Using various machine learning techniques (such as latent class analysis and cluster analysis), several population cohorts have deciphered various patterns of sensitization from childhood to adolescence and their differential risks for various allergic outcomes at later ages.^{6–10} While strong associations between AD and food sensitization are known, fewer studies have examined how manifestations of AD co-occur with these various sensitization patterns, especially at a younger age. One such study clearly showed that early AD, especially if associated with early food sensitization was highly associated with subsequent allergic disease, supporting the “atopic march” hypothesis.³ However, latent class analysis of data from the Manchester Asthma and Allergy Study (MAAS) suggested not all allergy leads to the atopic march, and that the atopic march may not be causal in subsequent development of allergic diseases.¹¹

The heterogeneity of methods used in past cross-sectional and cohort studies examining sensitization and AD as risk factors for allergic diseases makes interpretation difficult. Some examined only incident cases of AD,^{12,13} others assessed the risk of AD among all children and some did not distinguish between different types of sensitization.^{1,12–18} Ignoring the heterogeneity in the type of allergic sensitization (food, inhalant, food and inhalant, mono-sensitization and poly-sensitization), time of manifestation (persistent or transient) and AD status by simply categorizing children as “atopic” and “non-atopic” may obscure differential impacts of these risk factors.^{6–10} For instance, an increased risk for hospitalizations due to asthma was only found among children who were poly-sensitized at an early age but not among those sensitized at a later age.¹⁰ A recently reported unsupervised cluster analysis of data from the Copenhagen at-risk cohort from 6 months to 6 years of age identified 7 age- and allergen-specific patterns. Asthma was associated solely with the allergen pattern of dog/cat/horse, whereas AD was associated with all food and inhalant allergen clusters apart from house dust mite and allergic rhinitis with 5 of the 7 patterns.⁹

Machine learning techniques provide opportunities to explore the heterogeneity inherent in longitudinal patterns of development of symptoms of allergic diseases and of biological markers such as food and inhalant sensitization. We have applied latent class analysis to data from the Canadian Healthy Infant Longitudinal Development (CHILD) Study to examine the relationships between changes in AD and patterns of allergic sensitization from birth to age 3 years and the development of allergic diseases by that age, to better elucidate these patterns at an earlier age.

2 | MATERIALS AND METHODS

2.1 | Data collection

The CHILD Study recruited 3623 pregnant women from the general population across 4 Canadian provinces (Edmonton (Alberta), Toronto (Ontario), Vancouver (British Columbia) and Winnipeg (Manitoba). Excluding 83 children who were ineligible at birth and 45 families who failed to begin the study,¹⁹ we are following 3495 infants from pregnancy to age 5 years. The current analysis is based on 2629 children who completed skin prick tests (SPT) to common food and inhalant allergens at both 1 year and 3 years of age, and whose AD status was ascertained by clinical assessment by CHILD Study physicians at both ages. Food allergens tested included peanut, milk, egg white and soy. At 1 year and 3 years, inhalant allergens tested included *alternaria tenuis*, cat hair, dog epithelium, house dust mites (Der.pt and Der.f) and cockroach. Additional inhalants, namely *cladosporium*, *penicillium*, *aspergillus fumigatus*, trees, grass, weeds and ragweed were tested only at 3 years of age. At 3 months, 6 months, 1 year, 18 months, 2 years, 2.5 years and 3 years, parents were asked to complete questionnaires that inquired about the children’s skin rashes, allergic histories, medication and if the child had been diagnosed with eczema or atopic dermatitis by a health care professional during each time period. We accepted the terms eczema and atopic dermatitis as equivalent. Parents were also asked to report their own allergic history and completed similar skin prick tests (SPT). Full details on recruitment and procedures can be found in previous publications.¹⁹

2.2 | Definitions

Atopic status was determined by wheal size from SPTs, calculated as the average of the diameter and its perpendicular measured at the mid-point of the maximum length, from which was subtracted any wheal size produced by a negative control (glycerine) (Appendix S1). A child or parent was considered atopic if they produced a wheal ≥ 2 mm to at least one allergen.²⁰ In the very rare cases ($n = 3$) where skin testing to a specific allergen was declined due to previous reactions, external test results when available were used to determine sensitization status.

To reduce the child’s apprehension and because infants’ arms were small, some children had skin tests performed on the back rather than on the arms. Mean histamine wheal sizes on the back were consistently larger at both ages necessitating application of a correction factor of 0.82 to all wheals measured on the back (see details in Appendix S1).

During the clinic visits at 1 and 3 years, children were assessed for AD, asthma, allergic rhinitis and food allergy by interviewing parents about symptoms in the past year. Interviews were conducted by CHILD Study physicians (ABB, PJM, PS, SET; all experienced paediatricians specializing in allergy and asthma) or highly trained health care professionals under the guidance of these CHILD Study

physicians. Diagnoses were recorded as “definite,” “possible” or “no” for each allergic condition. In the current analysis, only “definite” cases were considered positive. The methods by which these diagnoses were established can be found in Appendix S2.

To ensure that the diagnosis of AD by 1 year of age was inclusive, we combined multiple sources of reporting.¹⁴ Based on the questionnaires, itchy wet or red rashes reported on the face, inside of elbows, wrist/hands or back of knees which were treated with corticosteroid creams were considered as definite AD. A child was also considered to have definite AD if parents reported a diagnosis made by a health care professional. Finally, definite AD was also recorded if the child was so diagnosed by our experienced study personnel during the clinical assessment visits, based on criteria from the UK Working Party.²¹ This required that children must have an itchy skin condition, together with one of the following three criteria: a history of involvement of the skin creases of elbows, behind the knees, front of ankles or around the neck; a history of general dry skin in the last year; or visible flexural eczema or eczema involving the cheeks/foreheads and outer limbs.

A parental history of allergic diseases was defined as at least one parent reporting a history of food allergies, eczema, asthma or allergic rhinitis (or hayfever). More details on definitions of variables can be found in Appendix S3.

2.3 | Statistical analysis

2.3.1 | Latent class analysis

First, we examined the changes in sensitization patterns from age 1 to 3 years for each allergen tested at both ages using McNemar's test for paired data. To understand the overall patterns of sensitization and AD within the data, we employed an exploratory, unsupervised latent class analysis (LCA), with no preconceived hypothesis. Ten factors that were assessed at 1 and 3 years of age were evaluated to reveal distinct patterns of sensitization and AD. These factors were as follows: AD at 1 and 3 years, food sensitization at 1 and 3 years, inhalant sensitization at 1 and 3 years, poly-sensitization to foods at 1 and 3 years and poly-sensitization to inhalants at 1 year and 3 years. In performing the LCA, we considered models ranging from 1 to 6 classes; we selected the optimum number of classes using the Bayesian information criterion (BIC) and Akaike information criterion (AIC), as well as considering the model that was most interpretable.^{22,23} The model then generated a posterior probability of membership in each identified class for each child. Subsequently, each child was assigned to the class that yielded the highest posterior probability among all the available classes. The “poLCA”²⁴ package in R 3.3.2²⁵ was used to perform the LCA.

2.3.2 | Risk for allergic diseases due to trajectories in sensitization patterns and AD

Following results from our LCA model, the identified groups were assessed for differences in the following sociodemographic factors:

sex, parental sensitization (SPT-based), parental history of allergic diseases, pet ownership, ethnicity, smoking in the house, breastfeeding, presence of older siblings, household income, parental education and study centre using a chi-square test (or Fisher's exact test where appropriate).^{12,13} To account for uncertainties in group assignment based on LCA, frequencies were weighted by the child's posterior probability of belonging to each group, rather than directly counting the number of children belonging in each group.^{26,27} These weights were calculated and rounded to the nearest whole number.²⁸ Demographic variables garnering significant chi-square results were followed with a post hoc comparison with the reference group (“healthy”); *P*-values were adjusted using the Holm-Bonferroni correction.²⁹

Finally, we evaluated the risks for demonstrating allergic diseases (allergic rhinitis, food allergy and asthma) at 3 years for each of the identified LCA groups compared to the “healthy” group. Odds ratios were obtained from multivariable logistic regression models with robust variance to account for the strong clustering effects of study centre.³⁰ To reduce bias, regression models were also weighted for each individual probability of group assignment.^{26,27} For example, a child with a 0.7 probability of belonging to group 1 and 0.3 probability of belonging to group 2 was considered as two observations in the regression model, contributing to the count in both groups, and weighted accordingly. Sensitivity analysis was conducted to determine whether results differed when these regression weights were not accounted for and each child was only assigned to the group with the highest membership probability.

Using a more relaxed threshold of $P < .2$, demographic factors that were associated with both outcomes and predictor variables were included as adjustment variables, while ensuring that enough events were observed given the number of covariates.³¹ To calculate the robust variance while accounting for the regression weights, the software STATA 13 was used for this analysis.³²

After groups were determined, we conducted post hoc analyses to determine whether an increased risk for food allergy might be explained through food avoidance by children who were food sensitized at age 1 year.

3 | RESULTS

3.1 | Study population

Of the 2629 children eligible for this analysis, the majority had a white Caucasian mother (74.3%) with a tertiary education (58.3%) and middle to upper socio-economic status (57.1%). The majority of parents reported a history of ever having allergies and atopic diseases (77.0%), and more than half (57.5%) of mothers tested positive to at least one allergen (Table 1). There were no significant differences in the characteristics of children included in the study to the full cohort (Appendix S4). However, among those excluded from this analysis, we noted a lower prevalence of AD at 1 year, fewer fathers with asthma and fewer mothers with AD.

TABLE 1 Demographic of participants

Demographics	N (%) = 2629	
Sex		
Female	1232 (46.9)	
Male	1397 (53.1)	
Study centre		
Edmonton	560 (21.2)	
Toronto	544 (20.7)	
Vancouver	620 (23.6)	
Manitoba	905 (34.4)	
Parental information ^a		
Family income		
	N (%)	
\$0-49 999	311 (11.9)	
\$50 000-99,999	828 (31.6)	
\$100 000-149 999	667 (25.5)	
>\$150 000	563 (21.5)	
Prefer not say	248 (9.5)	
	Mother	Father
Any parental history (ever)		
Eczema or Atopic Dermatitis	1606 (62.4)	1020 (45.6)
Allergic Rhinitis (Hayfever)	1281 (49.7)	1082 (48.3)
Asthma	577 (22.4)	449 (20.1)
Food Allergies	571 (22.3)	387 (17.5)
Any history of allergic disease	1986 (77.0)	1525 (68.1)
Sensitized (≥ 1 positive skin test)	1496 (57.5)	1422 (67.9)
Parental ethnicity		
First Nation	101 (3.9)	101 (3.8)
South East Asian	334 (12.8)	265 (10.1)
South Asian	75 (2.9)	93 (3.6)
Black	48 (1.8)	80 (3.1)
White	1941 (74.3)	1958 (74.8)
Other	112 (4.3)	108 (4.1)
Unknown	2 (0.1)	14 (0.5)
Highest parental education		
Graduate degree or higher	486 (19.1)	393 (15.5)
Postsecondary graduation	1485 (58.2)	1364 (53.9)
Some postsecondary	376 (14.7)	412 (16.3)
Secondary graduation (no postsecondary) or less	203 (8.0)	364 (14.4)

^aMay not sum to 2629 due to missing data.

3.2 | Prevalence of AD and allergic sensitization

At one year, 20.0% of children had AD and 13.5% were sensitized to at least one allergen; at 3 years, 22.1% had AD and 14.4% were sensitized. The majority of allergic sensitization at 1 year was to a single food; very few children were sensitized to inhalants. Conversely, at 3 years, more children were sensitized to inhalants; many were poly-sensitized and fewer children were sensitized to food (Table 2).

TABLE 2 Pattern of sensitizations and atopic dermatitis at 1 year and 3 years old

N (%) = 2629	1 Year	3 Year
Any food sensitization	284 (10.8)	155 (5.9)
Peanut	130 (5.0)	106 (4.0)
Cow's milk	47 (1.8)	33 (1.3)
Egg white	190 (7.2)	66 (2.5)
Soybean	21 (0.8)	14 (0.5)
Any inhalant sensitization	104 (4.0)	305 (11.6)
Alternaria Tenuis	19 (0.7)	16 (0.6)
Cat Hair	44 (1.7)	133 (5.1)
Dog Epithelium	19 (0.7)	29 (1.1)
D. Pteronyssinus	11 (0.4)	41 (1.6)
D. Farinae	14 (0.5)	42 (1.6)
Cockroach	16 (0.6)	27 (1.0)
Cladosporium	-	29 (1.1)
Penicillium mixed	-	9 (0.3)
Aspergillus fumigatus	-	13 (0.5)
Trees Midwest	-	42 (1.6)
Grass mix	-	53 (2.0)
Weeds	-	26 (1.0)
Ragweed mixed	-	22 (0.8)
Any sensitization (food or inhalant)	355 (13.5)	378 (14.4)
Poly-sensitization (food or inhalant)	109 (4.2)	166 (6.3)
Poly-sensitization to food	81 (3.1)	54 (2.1)
Poly-sensitization to inhalant	15 (0.6)	105 (4.0)
Number of positive skin tests (≥ 2 mm wheal)		
None	2274 (86.5)	2251 (85.6)
1 allergen	246 (9.4)	212 (8.1)
2 allergens	74 (2.8)	88 (3.4)
3 or more allergens	35 (1.3)	78 (3.0)
Reported or diagnosed with AD or eczema	527 (20.0)	582 (22.1)

The most common food sensitization at one year was to egg white, which decreased significantly at age 3 (7.2% to 2.5%, $P < .001$). At 3 years, peanut allergy was the most common food sensitization, although the rate had decreased significantly since age 1 year (4.9% to 3.9%; $P < .04$). Sensitization to cow milk also decreased significantly between age 1 and 3 years (1.8% to 1.1%, $P < .03$) but not soybean, which was the least common food sensitization at both ages (0.8% to 0.5%, $P < .29$).

The most common inhalant sensitization at both ages was to cat hair, which also demonstrated the largest increase with age (1.7% to 5.0%, $P < .001$). Sensitizations to both house dust mites were also common at both ages and increased significantly at age 3 (Der.pt: 0.4% to 1.5%, $P < .001$; Der.f: 0.6% to 1.6%; $P < .001$). Sensitization to alternaria tenuis was the least common at age 3 with no discernible changes between the two ages (0.7% to 0.6%; $P = .61$). Between age 1 and 3 years, more children were sensitized to dog epithelium (0.7% to

1.1%; $P = .11$) and cockroach (0.6% to 1.0%, $P = .09$); these differences were approaching significance. The changes in rates of sensitization for each allergen are shown in Figure 1.

3.3 | Trajectory of sensitization patterns and AD from LCA

A five-class structure yielded the lowest BIC among all models considered; a six-class structure gave a slightly smaller AIC; however, this difference was negligible (Appendix S5). After considering the interpretability of 4, 5 and 6 classes models, as well as recognizing BIC as the preferred model diagnostic criteria,²³ a 5-class structure was selected as the optimum model; there was an excellent separation of classes, with entropy of 0.91.³³ These 5 classes were as follows:

Healthy: Of all eligible children, 81.8% clustered into a “healthy” group; these children demonstrated very little AD or sensitization at either age.

Atopic Dermatitis: The group accounts for 7.6% of the children. This group displayed very little sensitization at either age, but at both ages, a large number had AD (1Y: 100%; 3Y: 85.2%).

Inhalant sensitization: The group accounts for 3.5% of children in the study. This group was characterized by little sensitization of any type at age 1 but substantial inhalant sensitization at age 3 years. They demonstrated a moderate level of AD at both ages (1Y: 30.0%; 3Y: 41.3%). Sensitization at age 1 tended to be mono-sensitization, whereas the surge of inhalant sensitization at age 3 was primarily poly-sensitization (65.0%); food sensitization remained low at age 3.

Transient sensitization: This group accounts for 4.1% of the children. They displayed a unique pattern where 100% were sensitized to food with few inhalant sensitizations at age 1 year; as they aged, food sensitization decreased significantly (to 30.8%) and inhalant sensitization remained low (11.4%). Food

sensitization at age 1 was primarily mono-sensitization (74.0%), and the level of AD was moderate (1Y: 28.7%; 3Y: 38.4%). As most did not retain sensitization at age 3, this was termed the “transient sensitization” group.

Persistent sensitization: The last group identified was the smallest, accounting for 3.2% of children. Almost all in this group (87.5%) were food sensitized at age 1 year, with little inhalant sensitization, but unlike the “transient sensitization” group, 100% were food sensitized at age 3; they were also largely poly-sensitized to food at both ages (1Y: 56.7%; 3Y: 62.8%). This group was characterized by higher prevalence of AD at both ages compared to other groups (1Y: 77.1%, 3Y: 67.6%). Many of them also acquired inhalant sensitization at age 3 years (72.2%), primarily mono-sensitization (61.0%). Given the persistence of their food sensitizations, this group is referred to as “persistent sensitization.” Figure 2 illustrates the probability distribution of the 10 key variables within the identified LCA groups (more details in Appendix S6).

3.4 | Demographic factors related to latent class group memberships

Sex, parental sensitization, pet ownership at 1 year, ethnicity and study centre were all significantly associated with the patterns of allergic sensitization and AD that emerged from the LCA (overall $P < .01$). However, having an older sibling, reported allergies in parents, income, smoking in the home, any breastfeeding at 1 year and parental education were not significantly associated with the patterns of AD and sensitization (Table 3).

Post hoc analysis revealed the direction of these observed differences. Males were more likely to be in the “inhalant sensitization” group ($P = .02$) and especially the “persistent sensitization” group ($P = .02$), with twice as many males as females in this group. Similarly, a strong difference in parental sensitization was found in the “persistent sensitization” group, where 90.7% of children had at least one parent with a




FIGURE 1 Changes in sensitization patterns from 1 year to 3 year*.

*P-values were obtained by McNemar's test of proportions.

positive skin prick test ($P < .01$). Children in the persistent sensitization group were less likely to have lived with pets by age one year compared to children in the healthy group ($P = .03$). Compared to the “healthy” group, children in the other four groups were more likely to be non-white ($P < .05$ for all). Although the cohort sample’s composition was primarily white Caucasian, 66.3% of those in the “persistent sensitization” group were non-white, indicating a strong effect of ethnicity ($P < .001$). Compared to healthy children, a higher proportion of “persistent sensitization” children came from Edmonton and Vancouver, while Winnipeg had a lowest proportion of “persistent sensitization” children. On the other hand, children living in Edmonton and Winnipeg represented the smallest proportion in the “inhalant sensitization” group, whereas the children living in Vancouver represented a higher proportion of this group.

After considering the association with allergic diseases presenting at 3 years, the following factors were retained as potential confounders for subsequent analyses: sex, parental sensitization, reported parental allergies, any breastfeeding, ethnicity and any pets.

3.5 | Risks for allergic diseases at 3 years by latent class memberships

3.5.1 | Asthma

Compared to healthy children, children in the “persistent sensitization” group had the greatest risk of having asthma at 3 years,

with an almost 12-fold increased risk (aOR: 11.44; 95% CI: 5.60, 23.37). Risk was also increased among children in the “transient sensitization” group despite no longer being sensitized at 3 years of age, as well as in the “inhalant sensitization” group; albeit risks in these groups were much lower than the “persistent sensitization” group (aOR: 3.84, 95% CI: 2.94, 5.00; aOR: 2.90, 95% CI: 1.45, 5.78, respectively). The risk of having asthma at 3 years among children in the “atopic dermatitis” group was also increased, albeit the lowest among all other groups (aOR: 1.65; 95% CI: 1.05, 2.57). When groups were not weighted according to their posterior probability, the risk for asthma in the “atopic dermatitis” group was no longer significant (Appendix S7).

3.5.2 | Allergic rhinitis

Children in the “persistent sensitization” group were six times more likely to develop allergic rhinitis at age 3 compared to children in the “healthy” group (aOR: 6.37, 95% CI: 2.56, 15.84). However, unlike asthma, the odds for developing rhinitis among children in the “inhalant sensitization” group were just as high as in the “persistent sensitization” group (aOR: 7.45; 95% CI: 3.98, 13.94). Children in the “transient sensitization” and “atopic dermatitis” groups were also at increased but somewhat lower risk of developing rhinitis, (aOR: 2.71; 95% CI: 1.26, 5.83; aOR: 2.36; 95% CI: 2.13, 2.62, respectively).




FIGURE 2 Probability of atopic dermatitis and allergic sensitization conditional on latent class memberships

*Actual frequencies for each group and diagnostic of model fits from LCA can be found in Appendix S6.

■ Probability of exhibiting the associated characteristics.

TABLE 3 Association between demographic characteristics with patterns of sensitization and atopic dermatitis (N = 2629)

Groups (from LCA)	N (row %) ^a		P-value ^b <.001
	Sex		
	Females	Males	
Healthy	1037 (48.7)	1094 (51.3)	-
Atopic dermatitis	80 (40.8)	116 (59.2)	.07
Inhalant sensitization	38 (34.9)	71 (65.1)	.02
Transient sensitization	49 (45.8)	58 (54.2)	.62
Persistent sensitization	28 (32.6)	58 (67.4)	.02
χ^2	19.1		
Groups (from LCA)	Parental sensitization (SPT)		<.001
	No	Yes	
	Healthy	471 (22.2)	
Atopic dermatitis	32 (16.3)	164 (83.7)	.08
Inhalant sensitization	15 (13.9)	93 (86.1)	.08
Transient sensitization	13 (12.1)	94 (87.9)	.03
Persistent sensitization	8 (9.3)	78 (90.7)	.01
χ^2	19.8		
Groups (from LCA)	Parental reported allergic disease		.52
	No	Yes	
	Healthy	216 (10.3)	
Atopic dermatitis	14 (7.1)	183 (92.9)	-
Inhalant sensitization	9 (8.4)	98 (91.6)	-
Transient sensitization	8 (7.5)	98 (92.5)	-
Persistent sensitization	7 (8.2)	79 (91.9)	-
χ^2	3.2		
Groups (from LCA)	Pet ownership at 1 year		.02
	No	Yes	
	Healthy	1065 (50.3)	
Atopic dermatitis	104 (54.4)	91 (45.6)	.54
Inhalant sensitization	55 (59.3)	36 (40.7)	.23
Transient sensitization	63 (56.2)	45 (43.8)	.54
Persistent sensitization	56 (65.1)	29 (34.9)	.03
χ^2	11.7		
Groups (from LCA)	Ethnicity		<.001
	Non-White	White	
	Healthy	688 (32.4)	
Atopic dermatitis	85 (43.1)	112 (56.9)	.008
Inhalant sensitization	49 (45.4)	59 (54.6)	.01
Transient sensitization	48 (45.3)	58 (54.7)	.01
Persistent sensitization	57 (66.3)	29 (33.7)	<.001
χ^2	58.7		
Groups (from LCA)	Smoking in the home 1 Y		.32
	No	Yes	
	Healthy	1718 (81.2)	
Atopic dermatitis	162 (83.9)	31 (16.1)	-

(Continues)

TABLE 3 (Continued)

		Smoking in the home 1 Y				
Groups (from LCA)	No	Yes				
Inhalant sensitization	94 (86.2)	15 (13.8)				
Transient sensitization	91 (86.7)	14 (13.3)				
Persistent sensitization	73 (84.8)	13 (15.1)				
χ^2	4.7					
		Any breastfeeding at 1 Y				
Groups (from LCA)	No	Yes				
Healthy	1028 (51.2)	978 (48.8)				
Atopic dermatitis	116 (60.7)	75 (39.3)				
Inhalant sensitization	52 (51.5)	49 (48.5)				
Transient sensitization	48 (49.0)	50 (51.0)				
Persistent sensitization	39 (48.8)	41 (51.3)				
χ^2	7.0					
		Older sibling				
Groups (from LCA)	No	Yes				
Healthy	1403 (65.9)	725 (34.1)				
Atopic dermatitis	139 (70.6)	58 (29.4)				
Inhalant sensitization	75 (68.8)	34 (31.2)				
Transient sensitization	71 (67.0)	35 (33.0)				
Persistent sensitization	60 (69.8)	26 (30.2)				
χ^2	2.4					
		Study centre				
Groups (from LCA)	Edmonton	Toronto	Vancouver	Winnipeg	<.001	
Healthy	452 (21.2)	420 (19.7)	477 (22.4)	782 (36.7)	-	
Atopic dermatitis	38 (19.3)	48 (24.6)	47 (23.9)	64 (32.5)	.35	
Inhalant sensitization	14 (16.8)	31 (29.0)	41 (38.3)	17 (15.9)	<.001	
Transient sensitization	27 (25.5)	29 (27.4)	28 (26.4)	22 (20.8)	.02	
Persistent sensitization	25 (29.1)	15 (17.4)	27 (31.4)	19 (22.1)	.03	
χ^2	50.6					
		Income				
Groups (from LCA)	\$0-49 999	\$50 000-99 999	\$100 000-149 999	>\$150 000	Prefer not say	.59
Healthy	267 (12.6)	675 (31.8)	525 (24.8)	456 (21.5)	197 (9.3)	-
Atopic dermatitis	17 (8.6)	68 (34.5)	57 (28.9)	36 (18.3)	19 (9.6)	-
Inhalant sensitization	11 (10.0)	30 (27.5)	32 (29.4)	28 (25.7)	8 (7.3)	-
Transient sensitization	9 (8.5)	28 (26.4)	31 (29.2)	25 (23.6)	13 (12.3)	-
Persistent sensitization	8 (8.1)	28 (32.6)	22 (25.6)	18 (20.9)	11 (12.8)	-
χ^2	14.2					
		Highest education of either parents				
Groups (from LCA)	Secondary graduation or less	Some postsecondary	Postsecondary graduation	Graduate degree or higher	.53	
Healthy	121 (5.8)	210 (10.1)	1197 (57.6)	549 (26.4)	-	
Atopic dermatitis	8 (4.1)	19 (9.8)	128 (64.4)	42 (21.6)	-	
Inhalant sensitization	2 (2.8)	11 (10.2)	60 (55.5)	34 (31.5)	-	
Transient sensitization	3 (2.8)	11 (10.5)	64 (61.0)	27 (25.7)	-	

(Continues)

TABLE 3 (Continued)

Groups (from LCA)	Highest education of either parents				.53
	Secondary graduation or less	Some postsecondary	Postsecondary graduation	Graduate degree or higher	
Persistent sensitization	3 (3.7)	5 (6.1)	48 (58.5)	26 (31.7)	-
χ^2	11.0				

^aFrequencies are weighted for the posterior probabilities of each child belonging in each group.

^bAll pairwise p-values were presented in comparison with the "healthy." Pairwise comparisons were conducted only when an overall significance was detected. Pairwise comparison p-values were adjusted using the Fifer package in R.

χ^2 are the chi-square values for each categorical variable.

3.5.3 | Food allergy

The risk of having reported food allergy was most elevated for those in the "persistent sensitization" group (aOR: 110.74, 95% CI: 38.33, 319.97). Although the "transient sensitization" children were no longer exhibiting food sensitization at 3 years, they were still at high risk of having food allergy at 3 years compared with healthy children (aOR: 22.35, 95% CI: 13.06, 38.24). In spite of low rates of food sensitization at ages 1 and 3, children in the "atopic dermatitis" and "inhalant sensitization" groups also exhibited increased risks for developing reported food allergies compared to healthy children (aOR: 5.16, 95% CI: 3.03, 8.78; aOR: 7.37, 95% CI: 3.50, 15.54, respectively). Additional post hoc analysis supported the possibility that the increased risk of food allergy among children in the "persistent sensitization" group may have been related to children avoiding foods to which they were sensitized at 1 year; children in the "persistent sensitization" group were more likely to have avoided common food allergens in the past year than any other groups, including "transient sensitization" (Table 4).

Adjusted and unadjusted risks for all group memberships are summarized in Table 5. Results from sensitivity analysis ignoring the posterior probability weighting of each group memberships can be found in Appendix S7.

4 | DISCUSSION

This study has identified almost 20% of children in a general Canadian population exhibiting four distinct patterns of allergic sensitization and AD, resulting in differential risks for important allergic outcomes at 3 years including asthma, allergic rhinitis and food

allergy. Children with persistent poly-sensitization to foods from 1 to 3 years of age concomitant with AD were at the highest risk for all the studied allergic diseases. Children with AD only were at risk for rhinitis and food allergy.^{16,17} On the other hand, although children who demonstrated mono-sensitization to foods at 1 year tended to have transient sensitization that disappeared by 3 years, they were still at an increased risk for allergic outcomes, especially reported food allergy. Finally, as well reported in past studies,^{1,4,34} some children who were non-sensitized at 1 year developed inhalant sensitization by 3 years. The risk of reported food allergy among these children was less than in other groups, although their risk of developing allergic rhinitis was equal to that of the "persistent sensitization" children.

The term "atopic march" has been used to refer to the progression of atopic dermatitis to asthma and allergic rhinitis. However, we and others have noted that "atopic dermatitis" is often a misnomer as quite frequently AD is diagnosed in the absence of allergic sensitization. In another publication from this population-based, longitudinal birth cohort, we reported that AD without evidence of allergic sensitization was not associated with an increased risk of subsequent asthma,³⁵ whereas the combination of AD and allergic sensitization at age 1 year was associated with increased risks for both asthma and food allergy at 3 years. The combined effect of diagnosed AD and allergic sensitization was greater than the sum of their individual effects on these risks.

Our results have confirmed past studies that found similar sensitization patterns among children and their associated risks with allergic diseases at age 3 years. Roduit and colleagues undertook latent class analysis in 1038 children in the PASTURE cohort, reporting that while both early transient and early persistent AD were associated with food allergy, persistent AD increased the risk of asthma

TABLE 4 Class memberships and food avoidance in the past year (between 2 and 3 years)

Group	N (% column) ^a					P-value
	Healthy	Atopic dermatitis	Inhalant sensitization	Transient sensitization	Persistent sensitization	
Consume milk	1833 (97.45)	181 (97.84)	70 (94.59)	92 (97.87)	62 (88.57)	<.001
Avoid milk	48 (2.55)	4 (2.16)	4 (5.41)	2 (2.13)	8 (11.43)	
Consume peanut	1760 (95.39)	169 (92.35)	65 (92.86)	71 (84.52)	29 (50.00)	<.001
Avoid peanut	85 (4.06)	14 (7.65)	5 (7.14)	13 (15.48)	29 (50.00)	
Consume egg	1816 (97.06)	176 (95.14)	73 (97.33)	72 (81.82)	44 (70.97)	<.001
Avoid egg	55 (2.94)	9 (4.86)	2 (2.67)	16 (18.18)	18 (29.03)	

^aDoes not sum to 2629 due to missing food avoidance data.

TABLE 5 Odds for asthma, allergic rhinitis and reported food allergies at age 3 years within each identified pattern

Asthma at 3 Y	OR (N = 2616) (No. of events = 89)	aOR^a (N = 2419) (No. of events = 82)
Healthy	1.00 (Ref)	1.00 (Ref)
Atopic dermatitis	2.03 (1.44, 2.87)	1.65 (1.05, 2.57)
Inhalant sensitization	2.54 (1.06, 6.09)	2.90 (1.46, 5.78)
Transient sensitization	3.58 (2.67, 4.80)	3.84 (2.94, 5.00)
Persistent sensitization	9.57 (4.24, 21.58)	11.44 (5.60, 23.37)
Allergic rhinitis at 3 Y	(No. of events = 63)	(No. of events = 56)
Healthy	1.00 (Ref)	1.00 (Ref)
Atopic dermatitis	2.32 (2.13, 2.54)	2.36 (2.13, 2.62)
Inhalant sensitization	7.59 (2.84, 20.27)	7.45 (3.98, 13.94)
Transient sensitization	3.00 (1.42, 6.36)	2.71 (1.26, 5.83)
Persistent sensitization	6.07 (2.71, 13.56)	6.37 (2.56, 15.84)
Reported food allergy at 3 Y	(No. of events = 112)	(No. of events = 104)
Healthy	1.00 (Ref)	1.00 (Ref)
Atopic dermatitis	5.70 (3.39, 9.57)	5.16 (3.03, 8.78)
Inhalant sensitization	7.27 (3.24, 16.31)	7.37 (3.50, 15.54)
Transient sensitization	22.42 (11.32, 44.38)	22.35 (13.06, 38.24)
Persistent sensitization	128.51 (36.8, 451.50)	110.74 (38.33, 319.97)

^aAdjusted for sex, parental sensitization, parental reported allergies, any breastfeeding at 1 year, ethnicity, sex and any pets. Confidence intervals take into account the clustering effects of study centres.

OR, unadjusted odds ratio; aOR, adjusted odds ratio. Observations differ due to missing data on outcomes or covariates.

almost threefold, food allergy sevenfold and allergic rhinitis fourfold.³⁶ Lee and colleagues reported that early presentation of AD, associated with a high prevalence of atopy, was the strongest risk factor for subsequent asthma and increased airway responsiveness.³⁷ Similarly, Matricardi and colleagues found in the Multi-centre Allergy Study (MAS) that AD occurring before age 2 years in association with wheezing before age 3 years gave a positive predictive value of 41.4 for wheezing in early adolescence.³⁸ In the GUSTO cohort in Singapore, AD occurring before 18 months of age was associated with a threefold increased risk of new allergic sensitization; a similar threefold increase in risk of new sensitization was found for wheeze before 18 months.¹³

Strengths of the present study are its longitudinal design, large sample size and recruitment from a general population cohort that allows stronger inferences of causality for all children. Furthermore, the use of LCA allows us to examine the distinct effects of timing, sensitization types and AD on the development of allergic diseases, rather than simply combining effects of any sensitization and AD.

There are limitations to the study. First, misclassification may have occurred. There are no validated instruments for diagnosis of AD in children younger than 2 years of age.¹² We used the UK Working Party criteria,²¹ which may have underestimated the number of children with AD, as milder cases may not be diagnosed.¹² To address this, we included reported rashes that met accepted clinical

criteria for atopic dermatitis and were treated with corticosteroid creams. Similarly, food allergy was not identified by the gold standard of an oral food challenge, but rather relied on parent-reported symptoms of food allergy, which may be subject to bias. Secondly, we have not used specific IgE measurements as having some other studies, which may have identified sensitization to food or inhalant allergens over and above those detected by our panel of skin tests. Of necessity, we tested a limited number of allergens in determining the atopic status of the children. As reported by parents, some children reacted to other allergens not tested in this current study (such as kiwi, banana and sesame). For consistency, if these children did not have a positive reaction to the panel of allergens tested and did not provide external test results, they were classified as non-sensitized. Given that we tested the most common food allergens, we believe we have identified at least 90% of sensitized children. Finally, while our cohort was taken from the general population, there was an unintentional recruitment bias where the proportion of parents with a history of allergic diseases is somewhat higher than in the general population, which may limit generalizability.

The differences in food avoidance between “persistent sensitization” and “transient sensitization” are worth noting, in the light of recent literature indicating that food avoidance can exacerbate the risk for food allergy.^{39,40} Whenever mothers requested their children's skin test results at 1 year, we were careful to inform them that these results indicated food sensitization, not food allergy. However, our data do suggest children who were food sensitized at age 1 year were more likely to avoid common food allergens than the other groups. This may explain why children in the “transient sensitization” group were still at an increased risk of food allergy although their food sensitization had disappeared at age 3 years. Given the possibility of confounding by food avoidance, findings regarding food allergy in the current study should be interpreted with caution.

To our knowledge, this is the first general population-based prospective longitudinal birth cohort study that uniquely identifies trajectories of both AD and sensitization among 1- to 3-year-old children and assesses their effects on subsequent development of allergic diseases. These classes should be validated in an independent cohort, preferably with a more precise diagnosis of food allergy to assess the effect of transient sensitization on the development of food allergy. Further studies are required to identify interventions that could help prevent allergic diseases. Longitudinal data will also help determine whether these patterns persist, disappear or even re-emerge after remitting at 3 years, to confirm results from past retrospective or cross-sectional studies.^{4,34} The CHILD Study is currently collecting follow-up data at 5 years, which will allow us to examine this further.

In conclusion, our findings suggest that children with persistent food poly-sensitization with AD by age 3 years may require the most clinical attention to reduce the risk of subsequent development of allergic diseases. This study adds to the evidence of substantial heterogeneity within groups of “atopic” and “non-atopic” children.^{6,9,10} Researchers and clinicians need to be cognizant of the problems associated with labelling children simply as “atopic” and “non-atopic” without considering timing of their atopic history, type of sensitization and

AD status. These enhanced research practices will lead to more clinically relevant findings, which could assist in early identification of modifiable risk factors for acquiring allergic diseases.

ACKNOWLEDGEMENTS

We are grateful to all the families who took part in this study, and the whole CHILD team that includes interviewers, nurses, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers and receptionists. We are particularly grateful to Zihang Lu for statistical advice and Melanie St. John for meticulous data checking. The Canadian Institutes of Health Research (CIHR) and the Allergy, Genes and Environment (AllerGen) Network of Centres of Excellence (NCE) provided core funding for CHILD. Additional support has been provided by Health Canada, Environment Canada, Canada Mortgage and Housing Corporation, the Sick Children's Hospital Foundation, Don & Debbie Morrison, the Silver Thread Foundation and the Childhood Asthma Foundation. We also acknowledge the generosity of ALK-Abello, Mississauga, ON, Canada, in supplying all allergens for the study, and Lincoln Diagnostics Inc., Decatur, IL USA, for supplying the Duotip-Test II devices and skin testing kits. M Sears holds the AstraZeneca endowed chair in Respiratory Epidemiology.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ORCID

M. R. Sears  <http://orcid.org/0000-0002-7113-4917>

REFERENCES

- Chiu CY, Huang YL, Tsai MH, et al. Sensitization to food and inhalant allergens in relation to atopic diseases in early childhood: a birth cohort study. *PLoS ONE*. 2014;9:e102809.
- Kulig M, Bergmann R, Klettke U, Wahn V, Tacke U, Wahn U. Natural course of sensitization to food and inhalant allergens during the first 6 years of life. *J Allergy Clin Immunol*. 1999;103:1173-1179.
- Kjaer HF, Eller E, Andersen KE, Høst A, Bindslev-Jensen C. The association between early sensitization patterns and subsequent allergic disease. The DARC birth cohort study. *Pediatr Allergy Immunol*. 2009;20:726-734.
- de Jong AB, Dikkeschei L, Brand P. Sensitization patterns to food and inhalant allergens in childhood: a comparison of non-sensitized, monosensitized, and polysensitized children. *Pediatr Allergy Immunol*. 2011;22:166-171.
- Silvestri M, Rossi GA, Cozzani S, et al. Age-dependent tendency to become sensitized to other classes of aeroallergens in atopic asthmatic children. *Ann Allergy Asthma Immunol*. 1999;83:335-340.
- Custovic A, Sonntag H, Buchan I, Belgrave D, Simpson A, Prospero M. Evolution pathways of IgE responses to grass and mite allergens throughout childhood. *J Allergy Clin Immunol*. 2015;136:1645-1652.
- Lazic N, Roberts G, Custovic A, et al. Multiple atopy phenotypes and their associations with asthma: similar findings from two birth cohorts. *Allergy*. 2013;68:764-770.
- Stoltz D, Jackson D, Evans M, et al. Specific patterns of allergic sensitization in early childhood and asthma & rhinitis risk. *Clin Exp Allergy*. 2013;43:233-241.
- Schoos A-MM, Chawes BL, Melen E, et al. Sensitization trajectories in childhood revealed by using a cluster analysis. *J Allergy Clin Immunol* 2017. (in press); doi.org/10.1016/j.jaci.2017.01.041
- Simpson A, Tan VY, Winn J, et al. Beyond atopy: multiple patterns of sensitization in relation to asthma in a birth cohort study. *Am J Respir Crit Care Med*. 2010;181:1200-1206.
- Belgrave D, Granell R, Simpson A, et al. Developmental profiles of eczema, wheeze, and rhinitis: two population-based birth cohort studies. *PLoS Med*. 2014;11:e1001748.
- Lowe A, Abramson M, Hosking C, et al. The temporal sequence of allergic sensitization and onset of infantile eczema. *Clin Exp Allergy*. 2007;1:536-542.
- Loo E, Sim J, Goh A, et al. Predictors of allergen sensitization in Singapore children from birth to 3 years. *Allergy, Asthma Clin Immunol*. 2016;12:26.
- Martin PE, Eckert JK, Koplin JJ, et al. Which infants with eczema are at risk of food allergy? Results from a population-based cohort. *Clin Exp Allergy*. 2015;45:255-264.
- Ha EK, Baek JH, Lee S-Y, et al. Association of polysensitization, allergic multimorbidity, and allergy severity: a cross-sectional study of school children. *Int Arch Allergy Immunol*. 2016;171:251-260.
- Ziyab A, Karmaus W, Zhang H, et al. Association of filaggrin variants with asthma and rhinitis: is eczema or allergic sensitization an effect modifier? *Int Arch Allergy Immunol*. 2014;164:308-318.
- Marenholz I, Kerscher T, Bauerfeind A, et al. An interaction between filaggrin mutations and early food sensitization improves the prediction of childhood asthma. *J Allergy Clin Immunol*. 2009;123:911-916.
- Kurukulaaratchy R, Matthews S, Arshad S. Defining childhood atopic phenotypes to investigate the association of atopic sensitization with allergic disease. *Allergy*. 2005;60:1280-1286.
- Subbarao P, Anand S, Becker A, et al. The Canadian Healthy Infant Longitudinal Development (CHILD) Study: examining developmental origins of allergy and asthma. *Thorax* 2015;70:998-1000.
- Sears MR, Herbison GP, Holdaway MD, et al. The relative risks of sensitivity to grass pollen, house dust mite and cat dander. *Clin Exp Allergy*. 1989;19:419-424.
- Williams H, Jburney P, Pembroke A, Hay R. The UK Working Party's diagnostic criteria for atopic dermatitis. III. Independent hospital validation. *Br J Dermatol*. 1994;131:406-416.
- Garrett ES, Zeger SL. Latent class model diagnosis. *Biometrics*. 2000;56:1055-1067.
- Nylund KL, Asparouhov T, Muthén B. Deciding on the number of classes in latent class analysis and growth mixture modeling: a Monte Carlo simulation study. *Struct Equ Model*. 2007;14:535-569.
- Linzer DA, Lewis JB. polCA: an R package for polytomous variable latent class analysis. *J Stat Softw*. 2011;42:1-29.
- R Core Team. *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing; 2016.
- Savenije OE, Granell R, Caudri D, et al. Comparison of childhood wheezing phenotypes in 2 birth cohorts: ALSPAC and PIAMA. *J Allergy Clin Immunol*. 2011;127:1505-1512.
- Bray BC, Lanza ST, Tan X. Eliminating bias in classify-analyze approaches for latent class analysis. *Struct Equ Modeling*. 2015;22:1-11.
- Meindl B, Templ M, Alfons A, Kowarik A, Ribatet M. simPop: Simulation of Synthetic Populations for Survey Data Considering Auxiliary Information. R Packag version 0.60. 2017;
- Holm S. A simple sequentially rejective multiple test procedure. *Scand J Stat*. 1979;6:65-70.
- Williams RL. A note on robust variance estimation for cluster-correlated data. *Biometrics*. 2000;56:645-646.
- Vittinghoff E, McCulloch C. Relaxing the rule of ten events per variable in logistic and Cox regression. *Am J Epidemiol*. 2007;165:710-718.

32. StataCorp. *Stata Statistical Software: release 13*. College Station, TX: StataCorp; 2013.
33. Celeux G, Soromenho G. An entropy criterion for assessing the number of clusters in a mixture model. *J Classif*. 1996;13:195-212.
34. Fiocchi A, Pecora V, Petersson CJ, et al. Sensitization pattern to inhalant and food allergens in symptomatic children at first evaluation. *Ital J Pediatr*. 2015;41:96.
35. Tran MM, Lefebvre DL, Dharma C, et al. Predicting the atopic march: results from the Canadian Healthy Infant Longitudinal Development Study. *J Allergy Clin Immunol*. 2017. (in press).
36. Roduit C, Frei R, Depner M, et al. Phenotypes of atopic dermatitis depending on the timing of onset and progression in childhood. *JAMA Pediatr*. 2017;171:655-662.
37. Lee E, Lee S, Kwon J, et al. Atopic dermatitis phenotype with early onset and high serum IL-13 is linked to the new development of bronchial hyperresponsiveness in school children. *Allergy*. 2016;71:692-700.
38. Matricardi P, Illi S, Grüber C, et al. Wheezing in childhood: incidence, longitudinal patterns and factors predicting persistence. *Eur Respir J*. 2008;32:585-592.
39. Tran M, Lefebvre DL, Dai D, et al. Timing of food introduction and development of food sensitization in a prospective birth cohort. *Pediatr Allergy Immunol*. 2017;28:471-477.
40. Du Toit G, Roberts G, Sayre PH, et al. Randomized trial of peanut consumption in infants at risk for peanut allergy. *N Engl J Med*. 2015;372:803-813.

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article.

How to cite this article: Dharma C, Lefebvre DL, Tran MM, et al. Patterns of allergic sensitization and atopic dermatitis from 1 to 3 years: Effects on allergic diseases. *Clin Exp Allergy*. 2018;48:48-59. <https://doi.org/10.1111/cea.13063>

APPENDIX 1

CHILD STUDY INVESTIGATORS

Subbarao P (Director), The Hospital for Sick Children & University of Toronto; Turvey SE (co-Director), University of British Columbia; Anand SS, McMaster University; Azad MB, University of Manitoba; Becker AB, University of Manitoba; Befus AD, University of Alberta; Brauer M, University of British Columbia; Brook JR, University of Toronto; Chen E, Northwestern University, Chicago; Cyr MM, McMaster University; Daley D, University of British Columbia; Dell SD, The Hospital for Sick Children & University of Toronto; Denburg JA, McMaster University; Duan QL, Queen's University; Eiwegger T, The Hospital for Sick Children & University of Toronto; Grasmann H, The Hospital for Sick Children & University of Toronto; HayGlass K, University of Manitoba; Hegele RG, The Hospital for Sick Children & University of Toronto; Holness DL, University of Toronto; Hystad P, Oregon State University; Kobor M, University of British Columbia; Kollmann TR, University of British Columbia; Kozyrskyj AL, University of Alberta; Laprise C, Université du Québec à Chicoutimi; Lou WYW, University of Toronto; Macri J, McMaster University; Mandhane PJ, University of Alberta; Miller G, Northwestern University, Chicago; Moraes TJ, The Hospital for Sick Children & University of Toronto; Paré P, University of British Columbia; Ramsey C, University of Manitoba; Ratjen F, The Hospital for Sick Children & University of Toronto; Sandford A, University of British Columbia; Scott JA, University of Toronto; Scott J, University of Toronto; Sears MR, (Founding Director), McMaster University; Silverman F, University of Toronto; Simons E, University of Manitoba; Takaro T, Simon Fraser University; Tebbutt SJ, University of British Columbia; To T, The Hospital for Sick Children & University of Toronto;